



In Vivo Evidence of Structural Brain Asymmetry in Musicians

Gottfried Schlaug, Lutz Jancke, Yanxiong Huang, Helmuth Steinmetz

Science, New Series, Volume 267, Issue 5198 (Feb. 3, 1995), 699-701.

Stable URL:

<http://links.jstor.org/sici?sici=0036-8075%2819950203%293%3A267%3A5198%3C699%3AIVEOSB%3E2.0.CO%3B2-F>

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

Science is published by American Association for the Advancement of Science. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/aaas.html>.

Science

©1995 American Association for the Advancement of Science

JSTOR and the JSTOR logo are trademarks of JSTOR, and are Registered in the U.S. Patent and Trademark Office. For more information on JSTOR contact jstor-info@umich.edu.

©2002 JSTOR

- of a cytosolic protein 60 kD in size over a distance of 150 to 200 μm will be about 3 to 5 min. Simple diffusional signaling between the synapse and cell body could thus account for the delay in the potentiation observed in the present study. Active retrograde transport is known to take place at a rate of about 30 to 180 $\mu\text{m}/\text{min}$ [R. D. Allen, J. Metzuzals, I. Tasaki, S. T. Brady, S. P. Gilbert, *Science* **218**, 1127 (1985)], which would allow more rapid signaling than diffusion.
17. The increase in cytosolic Ca^{2+} was shown by direct Ca^{2+} imaging with the fluorescence dye Fluo-3 in separate cultures, through use of methods previously described [Y. Dan and M.-m. Poo, *Nature* **359**, 733 (1992)].
 18. The transection was performed in normal culture medium to facilitate resealing of the plasma membrane. The medium was replaced with Ca^{2+} -free solution [115 mM NaCl, 2 mM MgCl_2 , 2.5 mM KCl, 10 mM HEPES, 3 mM EGTA, and 0.1% BSA (pH 7.3)] 20 min after the neurite transection.
 19. S. Davis *et al.*, *Science* **253**, 59 (1991); N. Stahl *et al.*, *J. Biol. Chem.* **268**, 7628 (1993); S. Davis and G. D. Yancopoulos, *Curr. Opin. Neurobiol.* **3**, 20 (1993).
 20. N. Y. Ip *et al.*, *Cell* **69**, 1121 (1992).
 21. N. Stahl *et al.*, *Science* **263**, 92 (1994).
 22. C. Lütticken *et al.*, *ibid.*, p. 89 (1994); A. Bonni, D. A. Frank, C. Schindler, M. E. Greenberg, *ibid.*, **262**, 1575 (1993).

23. S. Saadat, M. Sendtner, H. Rohrer, *J. Cell. Biol.* **108**, 1807 (1989); C. Kalberg, S. Y. Yung, J. A. Kessler, *J. Neurochem.* **60**, 145 (1993).
24. Addition of actinomycin D at 2.5 $\mu\text{g}/\text{ml}$ to *Xenopus* embryonic cell cultures was shown to inhibit >95% of [^3H]uridine incorporation into the trichloroacetic acid-precipitable cell fraction from these cultures [D. K. O'Dowd, *Nature* **303**, 619 (1983)]. For blocking of RNA synthesis in *Xenopus* embryos, actinomycin D (25 $\mu\text{g}/\text{ml}$) has been used [G. Grafi and G. Gallii, *FEBS Lett.* **336**, 403 (1993)]. Incubation with cycloheximide at 0.6 μM (or 0.17 $\mu\text{g}/\text{ml}$) for 2 hours was found to achieve about 90% inhibition of [^3H]leucine incorporation into the acid-precipitable fraction of these *Xenopus* cultures [L. A. C. Blair, *J. Neurosci.* **3**, 1430 (1983)]. Higher concentrations of these drugs were used in the present study to ensure more complete inhibition.
25. M. Rende *et al.*, *Glia* **5**, 25 (1992); M. Sendtner, K. A. Stöckli, H. Thoenen, *J. Cell Biol.* **118**, 139 (1992); Friedman *et al.*, *Neuron* **9**, 295 (1992).
26. We thank Regeneron Inc. for providing CNTF and BDNF, L.-q. Liu for technical assistance, and E. Lu, R. Girard, and J. Alder for helpful comments on the manuscript. Supported by grants from the NSF (IBN 22106) and the Office of Naval Research (N00014-90-J1865).

16 August 1994; accepted 15 November 1994

In Vivo Evidence of Structural Brain Asymmetry in Musicians

Gottfried Schlaug,*† Lutz Jäncke, Yanxiong Huang, Helmuth Steinmetz*

Certain human talents, such as musical ability, have been associated with left-right differences in brain structure and function. In vivo magnetic resonance morphometry of the brain in musicians was used to measure the anatomical asymmetry of the planum temporale, a brain area containing auditory association cortex and previously shown to be a marker of structural and functional asymmetry. Musicians with perfect pitch revealed stronger leftward planum temporale asymmetry than nonmusicians or musicians without perfect pitch. The results indicate that outstanding musical ability is associated with increased leftward asymmetry of cortex subserving music-related functions.

A number of studies have demonstrated that the left hemisphere of the brain is dominant in the production and comprehension of language in the vast majority of persons (1). Similar attempts to localize musical functions have yielded conflicting data, mainly because studies of amusia—that is, impairment of musical skills as a result of cerebral lesions—have failed to reveal structural-functional maps similar to those of language organization (2). This situation has now changed with the introduction of positron emission tomography (PET) to measure regional cerebral blood

flow and metabolism during the processing of verbal and nonverbal stimuli. Whereas left hemispheric activation sites are seen during phonological, lexical, or semantic language task performance (3), right hemispheric preponderances are found for melodic and pitch perception, at least in musically naive subjects (4). However, process-

ing strategies may differ among individuals depending on prior musical experience (or giftedness), as suggested by PET experiments (5) and by behavioral (6) and neurophysiological (7) studies.

These proposed functional differences have only been related to anecdotal postmortem descriptions of gross anatomical differences in the brains of eminent musicians compared to nonmusicians as well as pronounced interhemispheric asymmetry mainly of temporal lobe structures (8). In an unselected postmortem sample that established an anatomical marker for cerebral asymmetry, the size of a well-defined portion of the posterior superior temporal gyrus, termed the planum temporale (PT), was larger on the left side in the majority of brains (9). Asymmetry of the PT has been increasingly accepted as a substrate of left hemisphere dominance for language-related auditory processing because (i) asymmetry of the PT first appears in higher primates, suggesting a relation with the evolution of language (10); (ii) the left PT coincides with the center of Wernicke's speech area as identified by lesion studies (11); (iii) macroscopic asymmetry of the PT correlates with cytoarchitectonic asymmetry of association cortices thought to play a role in higher order auditory processing (12); and (iv) asymmetry of the PT is correlated with handedness, with left-handers being anatomically more symmetrical (13).

Rightward deviation from the usual pattern of cerebral asymmetry may be associated with increased giftedness for talents for which the right hemisphere is assumed to be important (14). This proposed relation has been partially substantiated by connections between nonright-handedness, atypical visuospatial lateralization, spatial giftedness, and musical talent (15). We have used high-resolution in vivo magnetic resonance morphometry of the PT as an index of laterality in 30 healthy, right-handed professional musicians and compared the results with those from nonmusicians matched for age, sex, and handedness (16–18).

Table 1. Means (\pm SD) for age, degree of anatomical planum temporale asymmetry (δ PT), and size of left and right PT determined with in vivo magnetic resonance morphometry in healthy, right-handed musicians and nonmusicians.

Subjects	Age	δ PT†	PT size (mm ²)	
			Left	Right
Musicians (n = 30)	26 (4)	-0.36 (0.25)*	1063 (189)	750 (187)
Perfect pitch (n = 11)	27 (5)	-0.57 (0.21)**	1097 (202)	611 (105)
No perfect pitch (n = 19)	26 (4)	-0.23 (0.17)	1043 (183)	830 (178)
Nonmusicians (n = 30)	26 (3)	-0.23 (0.24)	896 (236)	736 (263)

G. Schlaug, Y. Huang, H. Steinmetz, Department of Neurology, Heinrich-Heine-Universität Düsseldorf, D-40225 Düsseldorf, Germany.

L. Jäncke, Institute of General Psychology, Section of Cybernetical Psychology and Psychobiology, Heinrich-Heine-Universität Düsseldorf, D-40225 Düsseldorf, Germany.

*To whom correspondence should be addressed.


†Present address: Department of Neurology, Beth Israel Hospital, Boston, MA 02215, USA.

†Negative values indicate leftward asymmetry of the PT (16).

* $P = 0.028$ compared to nonmusicians.

** $P < 0.001$ compared to musicians without perfect pitch (21).

Fig. 1. Brain surface projections of the right and left planum temporale (PT) in a musician with perfect pitch (upper) and a nonmusician (lower). (Left) Views from the right; (middle) views from above with the left brain hemispheres to the reader's right; (right) views from the left. The images were reconstructed from stacks of 128 contiguous sagittal magnetic resonance image slices where the PT had been highlighted on each slice. The δ PT values are -0.77 for the musician and -0.39 for the nonmusician (16).



We found that the PT was more lateralized to the left in musicians ($P = 0.028$). Possession of perfect pitch explained most of the variation in the degree of PT asymmetry among musicians ($P < 0.001$) (19–21). Musicians with perfect pitch showed stronger leftward PT asymmetry compared to other musicians, whereas musicians without perfect pitch did not differ from controls (Table 1 and Fig. 1).

Our finding of increased leftward PT asymmetry among musicians should be seen in the following context. First, PET has demonstrated that the posterior superior temporal region, including the PT, is involved in music perception (5). Second, in one postmortem myeloarchitectonic study of a musician with melody deafness after circumscribed brain injury, the lesion was centered on the left PT, sparing the primary auditory and inferior parietal cortex (22). Third, gross left-right asymmetry of the PT, as measured in our study, reflects cytoarchitectonic asymmetries of auditory association areas located on the PT (12). Thus, our morphometric findings in musicians may suggest that the functional capacity of cortex shown to subserve musical functions increases with leftward structural asymmetry of this neural system. This result lends anatomical support to behavioral and electrophysiological evidence of a difference in lateralization of musical processing between musicians and nonmusicians, with more left-lateralized representation in musicians (6, 7). Our data concur with the general concept that, because of time constraints of interhemispheric transfer, efficiency of neuronal assemblies is expected to increase with the number of elements clustered in one hemisphere (23). In fact, this principle may be the essence of hemispheric specialization (23).

Our study does not reveal the mechanism creating structural asymmetry. Leftward PT asymmetry usually appears in the human fetus between the 29th and

31st gestational week (24), so that prenatal factors are likely to play a role. Nevertheless, considering that the maturation of fiber tracts and intracortical neuropil, two presumed determinants of gyral shape (25), are still progressing by the age of seven (26), it remains uncertain whether gross anatomy may also be susceptible to some postnatal plastic change, such as in response to specific stimulation (20). Our study demonstrates that individual variability in cognitive performance can covary with features of external brain morphology.

REFERENCES AND NOTES

1. T. Rasmussen and B. Milner, *Ann. N.Y. Acad. Sci.* **299**, 355 (1977); P. Satz, *Science* **203**, 1131 (1979).
2. Since the 19th century, a number of investigators have ascribed amusia to left or right hemispheric dysfunction. Both the occurrence of amusia without aphasia and of aphasia without amusia have been described (double dissociation), suggesting at least partial independence of the neural structures subserving the respective functions. For overviews, see N. Wertheim, in *Handbook of Clinical Neurology*, P. J. Vinken and G. W. Bruyn, Eds. (Elsevier, New York, 1969), vol. 4, pp. 195–206; A. R. Damasio and H. Damasio, in *Music and the Brain*, M. Critchley and R. A. Henson, Eds. (Heinemann, London, 1977), pp. 141–155; and J. Sergent, *Trends Neurosci.* **16**, 168 (1993).
3. S. E. Petersen, P. T. Fox, M. I. Posner, M. Mintun, M. E. Raichle, *Nature* **331**, 585 (1988); S. Wise *et al.*, *Brain* **114**, 1803 (1991); J. F. Demonet, R. Wise, R. S. J. Frackowiak, *Hum. Brain Mapping* **1**, 39 (1993).
4. R. J. Zatorre, A. C. Evans, E. Meyer, A. Gjedde, *Science* **256**, 846 (1992); R. J. Zatorre, A. C. Evans, E. Meyer, *J. Neurosci.* **14**, 1908 (1994).
5. J. C. Mazziotta, M. E. Phelps, R. E. Carson, D. E. Kuhl, *Neurology* **32**, 921 (1982). In this study, subjects who later described themselves as having used "specific, highly organized" analytic approaches consistently displayed left greater than right temporoparietal activation during a tonal memory task. Right greater than left activation during the same task was found in individuals reporting no specific strategy.
6. T. G. Bever and R. J. Chiarello, *Science* **185**, 537 (1974). After comparing ear advantages for monaurally presented melodies in musicians (right-ear advantage) and nonmusicians (left-ear advantage), the authors concluded that "being musically sophisticated has real neurological concomitants, permitting the utilization of a different strategy of musical appre-

hension that calls on left hemisphere functions." For an opposing view see R. J. Zatorre, *Neuropsychologia* **17**, 607 (1979).

7. M. Hirshkowitz, J. Earle, B. Paley, *Neuropsychologia* **16**, 125 (1978); E. Altenmüller, *Eur. Arch. Psychiatry Neurol. Sci.* **235**, 342 (1986); M. Besson, F. Faits, J. Requin, *Neurosci. Lett.* **168**, 101 (1994).
8. For an overview, see A. Meyer, in *Music and the Brain*, M. Critchley and R. A. Henson, Eds. (Heinemann, London, 1977), pp. 255–281.
9. N. Geschwind and W. Levitsky, *Science* **161**, 186 (1968).
10. G. H. Yeni-Komshian and D. A. Benson, *ibid.* **192**, 387 (1976).
11. N. Geschwind, *ibid.* **170**, 940 (1970).
12. C. von Economo and L. Horn, *Z. Neurol. Psychiatry* **130**, 678 (1930); A. M. Galaburda, M. LeMay, T. L. Kemper, N. Geschwind, *Science* **199**, 852 (1978); A. M. Galaburda, F. Sanides, N. Geschwind, *Arch. Neurol.* **35**, 812 (1978).
13. H. Steinmetz, J. Volkman, L. Jäncke, H.-J. Freund, *Ann. Neurol.* **29**, 315 (1991); H. Steinmetz and A. M. Galaburda, *Read. Writ.* **3**, 329 (1991); L. Jäncke, G. Schlaug, Y. Huang, H. Steinmetz, *NeuroReport* **5**, 1161 (1994); H. Steinmetz, A. Herzog, G. Schlaug, Y. Huang, L. Jäncke, *Cereb. Cortex*, in press.
14. N. Geschwind and A. M. Galaburda, *Arch. Neurol.* **42**, 428 (1985); *ibid.*, p. 521; *Cerebral Lateralization: Biological Mechanisms, Associations, and Pathology* (MIT Press, Cambridge, MA, 1987).
15. M. Hassler and N. Birbaumer, *Dev. Neuropsychol.* **4**, 129 (1988); M. Hassler, *Brain Cogn.* **13**, 1 (1990); M. Hassler, *Int. J. Neurosci.* **68**, 145 (1993).
16. In vivo magnetic resonance morphometry was performed as described and validated previously in human cadaver brains [H. Steinmetz *et al.*, *Brain Lang.* **39**, 357 (1990)] through use of a Siemens 1.5 T magnet and a 22-min fast-low-angle-shot magnetic resonance sequence covering the entire brain, yielding a 1.00 mm by 1.00 mm by 1.17 mm image voxel size. Each dataset consisted of 128 contiguous, 1.17-mm-thick sagittal slices. For off-line morphometry, the datasets obtained from musicians and nonmusicians were mixed and coded. They were analyzed independently by two observers (G.S. and H.S.) blinded for musician or control, and right or left hemisphere. The curved length of the PT was measured on each slice. The anterior border of the PT was defined as the bottom of Heschl's sulcus, and the posterior border was defined as the transition of the posterior wall of the descending into the ascending ramus of the Sylvian fissure (73). By adding up the single-slice measurements multiplied by slice thickness, morphometry of the left (L) and right (R) PT provided an estimate of its entire folded cortical surface (in square millimeters). As in previous studies (73) [A. M. Galaburda, J. Corsiglia, G. Rosen, G. F. Sherman, *Neuropsychologia* **25**, 853 (1987)], the degree of asymmetry was expressed as the coefficient δ PT = $(R - L) / (0.5(R + L))$. The interobserver (Pearson) correlation coefficient for δ PT was 0.93 ($P < 0.001$) for the present sample.
17. The professional musicians were recruited through announcements in three well-known music schools in Germany, as well as through personal contacts. All were classical musicians who had either finished their education or were receiving formal training at a music school. No amateur musicians were included. All string players ($n = 14$) were keyboard players as well but had a preference for string instruments; the others were keyboard players ($n = 16$). The age-, sex-, and handedness-matched nonmusicians were recruited through announcements in the local medical school. Most of them were medical students or young faculty members in university hospitals. None of them had ever played a musical instrument or received formal musical training. All subjects gave informed consent.
18. Handedness was assessed with the 12-item questionnaire of M. Annett [*Br. J. Psychol.* **61**, 303 (1970)]. Consistent right-handedness was defined as performance of all 12 tasks with the right hand, with up to two "either" preferences being acceptable. Three male musicians and three male nonmusicians were nonconsistent righthanders; all other subjects were consistent righthanders. Also, non-

musicians and musicians did not differ in body length ($P > 0.05$).

19. No formal test for perfect pitch was applied. Subjects were asked if they would be able to sing any tone without a reference tone and if they could name any tone that was given to them without a reference tone. In those musicians who described themselves as having perfect pitch ($n = 11$), this information was confirmed by review of their music school examination records for pitch discrimination task performance.

20. In distinguishing between musicians with or without perfect pitch, we were guided by a study of D. Sergeant [*J. Res. Mus. Educ.* 17, 135 (1969)] investigating the possession of perfect pitch in a large sample of professional musicians. In this study almost all musicians who began training before the age of seven had perfect pitch, but almost none of those beginning after the age of 11. Similar results were also found by others suggesting that early exposure and disposition may underly this ability [A. Bachern, *J. Acoust. Soc. Am.* 27, 751 (1955); C. L. Krumhansl, *Annu. Rev. Psychol.* 42, 277 (1991); M. Klein, M. G. H. Coles, E. Donchin, *Science* 223, 1306 (1984)].

21. For statistical analysis, δ PVT values (16) were sub-

jected to a one-way analysis of variance (ANOVA) with three levels (musicians with or without perfect pitch, and controls). In this ANOVA two planned orthogonal (linear) contrasts were computed, comparing musicians versus controls and musicians with perfect pitch versus musicians without [R. E. Kirk, *Experimental Design: Procedures for the Behavioral Sciences* (Wadsworth, Belmont, CA, 1968), pp. 69-98]. The results were as follows: orthogonal contrast for musicians versus controls: [$F(1,57) = 5.12, P = 0.028$]; orthogonal contrast for musicians with perfect pitch versus musicians without: [$F(1,57) = 16.18, P < 0.001$]. Prior to the ANOVA, a Bartlett-Box-test for homogeneity of variances of δ PVT had revealed no deviation from the homogeneity assumption [$F(2,4840) = 1.25, P = 0.32$].

22. K. Kleist, *Sensorische Aphasien und Amusien auf myeloarchitektonischer Grundlage* (Thieme Verlag, Stuttgart, 1959), pp. 32-43. Similar cases of musicians with or without musical impairment lacked precise pathoanatomical verification (2) [A. R. Luria, L. S. Tsvetkova, D. S. Futer, *J. Neurol. Sci.* 2, 288 (1965); T. Alajouanine, *Brain* 71, 229 (1948)].

23. J. L. Ringo, R. W. Doty, S. Demeter, P. Y. Simard, *Cereb. Cortex* 4, 331 (1994). On the basis of esti-

mates of interhemispheric conduction times and on a neuronal network model, the authors hypothesized that "finely detailed, time-critical neuronal computations . . . would be performed more quickly via shorter and faster intrahemispheric circuits," and more generally that "unilateral clustering of related memories and skills would manifest itself as hemispheric specialization."

24. S. F. Witelson and W. Pallie, *Brain* 96, 641 (1973); J. A. Wada, R. Clarke, A. Hamm, *Arch. Neurol.* 32, 239 (1975); J. G. Chi, E. C. Dooling, F. H. Gilles, *ibid.* 34, 346 (1977).

25. W. Welker, in *Cerebral Cortex*, E. G. Jones and A. Peters, Eds. (Plenum, New York, 1990), vol. 8B, pp. 3-136.

26. P. I. Yakovlev and A.-R. Lecours, in *Regional Development of the Brain in Early Life*, A. Minkowski, Ed. (Blackwell, Oxford, 1967), pp. 3-70.

27. Supported by the Deutsche Forschungsgemeinschaft (SFB 194) and the Hermann und Lilly Schilling Stiftung (H.S.). We thank U. Mödder (Institute for Diagnostic Radiology, Heinrich-Heine-Universität Düsseldorf) for use of the magnetic resonance imaging facility.

6 September 1994; accepted 28 November 1994

TECHNICAL COMMENTS

Mammalian Vestibular Hair Cell Regeneration

Birds and mammals are born with a full complement of inner ear hair cells, which were thought to be irreversibly lost when damaged (1). It is now well known that birds have the capacity to regenerate hair cells in their auditory and vestibular organs after damage by acoustic trauma or ototoxic drugs (2) and that these new cells can mediate functional recovery (3). Recent studies by A. Forge *et al.* (4) and by M. E. Warchol *et al.* (5) suggest that the vestibular epithelium of the mature mammalian inner ear may also have the ability to produce new hair cells by renewed mitotic activity in response to aminoglycoside injury *in vivo* (4) and *in vitro* (5). However, these reports do not provide convincing evidence that the DNA labeling, seen at a low frequency *in vitro*, is the source of the apparent recovery of hair cell apical surfaces observed *in vivo*.

Our study was undertaken to determine if cell division can be shown to give rise to new hair cells in normal mature mammalian vestibular epithelium or during the first 6 weeks after aminoglycoside ototoxicity. Three groups of young mature albino Hartley guinea pigs were used. The experimental animals in each group were treated with a single transtympanic injection of the ototoxic aminoglycoside, gentamicin, in the left ear (6). Animals in each control group were given an identical volume of 0.9% saline. The first group of animals was killed after 1 to 16 weeks and used for light microscopic evaluation of damage produced in the sensory epithelium of the

utricle (7). The second group, killed after 1 to 16 weeks, was used for scanning electron microscopy (SEM) (8) in order to compare our results with those of Forge *et al.* (4). In animals of the third group, an osmotic pump filled with [3 H]thymidine was implanted under the skin of the back with its output leading to a cannula inserted into the perilymphatic space before treatment with aminoglycoside (9). These animals were killed after 1 to 16 weeks (10).

Hair cell damage and loss was evident in the light microscopic sections and SEM analyses of tissue from gentamicin-treated animals (Fig. 1). Experimental animals had fewer hair cells than controls, particularly in the striolar region. Other signs of damage observed by light microscopy of SEM included nuclear pyknosis, nuclear swelling, vacuolization, cytoplasmic extrusion, and

stereocilia fusion. The extent of damage was variable at all survival times. At 1 or 2 weeks after gentamicin treatment, hair cell injury was limited primarily to the striolar region in 10 of 16 animals examined by SEM. In three of the animals damage was observed over a larger area, extending from the striola toward the periphery of the organ. Complete destruction of the sensory hair cells was observed in the remaining three animals. Four weeks after gentamicin administration, one animal displayed hair cell damage extending out from the striolar region; in the other animal blebbing and fusion of stereocilia were seen over the entire surface of the sensory epithelium. In the animal killed 4 months after gentamicin, the surface of the utricle continued to show damaged stereocilia bundles throughout the entire sensory epithelium. The average length of the sensory epithelium and the linear support cell density remained constant between the control and experimental animals (Table 1) (11). However, the linear hair cell density was 51 to 85% lower in experimental animals than controls ($P < 0.001$).

Table 1. Results of treatment with gentamicin on guinea pig utricle: Length of sensory epithelium, hair cell density, and support cell density. Measurements are averages (\pm standard deviation).

Animal number	Treatment (weeks)	Sensory epithelium length ($\times 0.1$ mm)	Hair cell density (per 0.1 mm)	Support cell density (per 0.1 mm)
94-01	1	7.2 (± 2.0)	1.6 (± 0.7)	11.6 (± 4.6)
94-13	1	9.3 (± 2.1)	3.4 (± 0.8)	9.9 (± 3.1)
93-42	1	7.2 (± 1.6)	2.6 (± 1.4)	10.4 (± 1.5)
94-06	4	7.4 (± 1.5)	1.3 (± 1.5)	8.4 (± 1.5)
94-05	6	8.7 (± 2.0)	1.6 (± 0.6)	7.1 (± 1.6)
93-65	6	7.2 (± 1.9)	4.3 (± 0.8)	10.7 (± 2.8)
93-57	0*	7.4 (± 1.6)	8.7 (± 1.7)	12.6 (± 3.3)
93-38	6*	8.9 (± 2.1)	6.6 (± 1.9)	10.2 (± 1.2)

*Control group received no gentamicin.